


## Erratum to: Influence of two types of El Niños on the East Asian climate during boreal summer: a numerical study

Zesheng Chen<sup>1</sup> · Zhiping Wen<sup>1</sup> · Renguang Wu<sup>2</sup> · Ping Zhao<sup>3</sup> · Jie Cao<sup>4</sup>

Published online: 4 October 2017  
© Springer-Verlag GmbH Germany 2017

### Erratum to: *Clim Dyn* (2014) 43:469–481 DOI [10.1007/s00382-013-1943-1](https://doi.org/10.1007/s00382-013-1943-1)

In the article, we said that “The JJA Niño 3 SST index is defined by the mean SSTA averaged over the region (150°–90°W, 5°S–5°N, right solid box in Fig. 1). Similarly, the JJA Niño 4 SST index is defined using SST over the region (160°E–150°W, 5°S–5°N, left solid box in Fig. 1).” However, the labels “Niño 3” and “Niño 4” are located in the wrong position in the original Fig. 1, even though the Niño 3 and Niño 4 SST indices are correctly calculated in the study. Below is the new Fig. 1.



**Fig. 1** Representation of the regions employed to calculate the indices used in this study, including Niño3 SST index, Niño4 SST index and western north Pacific summer monsoon index

The online version of the original article can be found under doi:[10.1007/s00382-013-1943-1](https://doi.org/10.1007/s00382-013-1943-1).

✉ Zhiping Wen  
eeswzp@mail.sysu.edu.cn

- <sup>1</sup> Center for Monsoon and Environment Research/  
Department of Atmospheric Sciences, Sun Yat-sen  
University, No. 135, Xingangxi Road, Guangzhou 510275,  
People’s Republic of China
- <sup>2</sup> Institute of Space and Earth Information Science/Department  
of Physics, Chinese University of Hong Kong,  
Shatin, Hong Kong, People’s Republic of China
- <sup>3</sup> Chinese Academy of Meteorological Sciences, Beijing,  
People’s Republic of China
- <sup>4</sup> Department of Atmospheric Sciences, Yunnan University,  
Kunming, People’s Republic of China