



Correction to: Pre-post analysis of a social capital-based exercise adherence intervention for breast cancer survivors with moderate fatigue: a randomized controlled trial

Sue Kim¹ · Yun Hee Ko¹ · Yoonkyung Song² · Min Jae Kang² · Hyojin Lee¹ · Sung Hae Kim³ · Justin Y. Jeon^{2,4} · Young Up Cho⁵ · Gihong Yi⁶ · Jeehee Han¹

Published online: 16 October 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: Supportive Care in Cancer (2020) 28:5281–5289.

<https://doi.org/10.1007/s00520-020-05363-7>

Tables 1, 2, and 3 incorrect in the original article, the correct tables are shown below:

Below is also the correct ethical statement:

Ethics approval and consent to participate An institutional review board was obtained from Severance Hospital, Korea (IRB 4–2017–0164).

The online version of the original article can be found at <https://doi.org/10.1007/s00520-020-05363-7>

✉ Jeehee Han
hjihee1225@hanmail.net

Sue Kim
suekim@yuhs.ac

Yun Hee Ko
urekahee@naver.com

Yoonkyung Song
lapine1019@gmail.com

Min Jae Kang
lejo096@naver.com

Hyojin Lee
kailuabreeze91@naver.com

Sung Hae Kim
wrlrtk@gmail.com

Justin Y. Jeon
jjeon@yonsei.ac.kr

Young Up Cho
youngup@yuhs.ac

Gihong Yi
gihong@hallym.ac.kr

- ¹ Mo-Im Kim Nursing Research Institute, College of Nursing, Yonsei University, 50-1, Yonsei-ro, Seodaemun-gu, Seoul, South Korea
- ² Department of Sports Industry Studies, Yonsei University, 50-1, Yonsei-ro, Seodaemun-gu, Seoul, South Korea
- ³ Department of Nursing, College of Health, Welfare and Education, Tongmyong University, Busan, South Korea
- ⁴ Exercise Medicine Center for Diabetes and Cancer Patients, ICONS, Yonsei University, 50-1, Yonsei-ro, Seodaemun-gu, Seoul, South Korea
- ⁵ Formerly affiliated with Yonsei University Health System, 50-1, Yonsei-ro, Seodaemun-gu, Seoul, South Korea
- ⁶ Department of Sociology, Hallym University, 1, Hallymdaehak-gil, Chuncheon-si, Gangwon-do 24252, South Korea

Table 1 Demographics and clinical characteristics (*N* = 48)

		Int. (<i>n</i> = 23)	Cont. (<i>n</i> = 25)	χ^2/t (p)
		Mean \pm SD (range) N (%)	Mean \pm SD (range) N (%)	
Age(years)		49.91 \pm 7.62 (34–67)	48.48 \pm 6.75(33–62)	.690(.494)
	30–39	1(4.3)	2(8.0)	.751(.861)
	40–49	11(47.8)	13(52.0)	
	50–59	9(39.1)	9(36.0)	
	60–69	2(8.7)	1(4.0)	
Marital status	Married	16(69.6)	17(68.0)	.167(.765)
	Unmarried	7(30.4)	8(32.0)	
Income (10,000won)	<300	12(52.2)	11(44.0)	.321(.773)
	\geq 300	11(47.8)	14(56.0)	
Employment status	No	16(69.6)	14(56.0)	.941(.313)
	Yes	7(30.4)	11(44.0)	
Religion	No	7(30.4)	7(28.0)	.034(.552)
	Yes	16(69.6)	18(72.0)	
Economic burden	No	12(52.2)	17(68.0)	1.255(.377)
	Yes	11(47.8)	8(32.0)	
Education level	<Middle school	1(4.3)	1(4.0)	4.333(.115)
	\geq High school	22(95.7)	24(96.0)	
Parental status	No	5(21.7)	4(16.0)	.259(.719)
	Yes	18(78.3)	21(84.0)	
Stage	I	5(21.7)	7(28.0)	1.790(.617)
	II	13(56.5)	15(60.0)	
	III	5(21.7)	3(12.0)	
Surgery type	Mastectomy	4(17.4)	8(32.0)	.243 (.342)
	Lumpectomy	19(82.6)	17(68.0)	
Time since diagnosis	<1	8(34.8)	11(44.0)	2.408(.300)
	1–2	13(56.5)	9(36.0)	
	\geq 2	2(8.7)	5(20.0)	
Chemotherapy	None	0(0)	2(8.0)	1.920(.490)
	Past	23(100)	23(92.0)	
Radiation therapy	None	1(4.3)	3(12.0)	.918(.632)
	Past	21(91.3)	21(84.0)	
	Present	1(4.3)	1(4.0)	
Endocrine therapy	No	11(47.8)	12(48.0)	1.790(.617)
	Yes	12(52.2)	13(52.0)	
Target therapy	No	12(52.2)	20(80.0)	4.591(.101)
	Yes	10(43.5)	5(20.0)	
	Unknown	1(4.3)	0(0.0)	
Triple-negative status	No	16(69.6)	14(56.0)	1.122(.571)
	Yes	6(26.1)	7(28.0)	
	Unknown	1(4.3)	4(16.0)	

Note. Int = Intervention group; Cont = Control group

Table 2 Homogeneity of dependent variables between groups (N = 48)

Variables	Int. (n = 23) Mean ± SD (possible range)	Cont. (n = 25) Mean ± SD (possible range)	t(p)
Fatigue	5.24 ± 1.53 (0–10)	5.37 ± 1.80 (0–10)	−0.268(.790)
Behavioral/Severity	5.55 ± 1.42 (0–10)	5.46 ± 2.26 (0–10)	0.181(.858)
Affective	5.84 ± 2.11 (0–10)	5.89 ± 2.55 (0–10)	−0.062(.951)
Sensory	4.70 ± 2.31 (0–10)	5.53 ± 1.87 (0–10)	−1.358(.181)
Cognitive/mood	4.80 ± 2.00 (0–10)	4.72 ± 2.05 (0–10)	0.834(.892)
Quality of life	85.58 ± 11.86 (0–148)	86.80 ± 20.31 (0–148)	−0.258(.798)
Physical wellbeing	17.73 ± 4.70 (0–28)	17.12 ± 6.48 (0–28)	0.376(.709)
Social/family wellbeing	15.88 ± 5.46 (0–28)	17.76 ± 5.20 (0–28)	−1.218(.230)
Emotional wellbeing	15.26 ± 3.55 (0–24)	15.36 ± 4.59 (0–24)	−0.083(.934)
Functional wellbeing	14.86 ± 4.54 (0–28)	16.28 ± 4.79 (0–28)	−1.044(.302)
Breast cancer subscale	21.82 ± 4.31 (0–40)	20.28 ± 5.38 (0–40)	1.089(.278)
Physical activity (MET)	2610.43 ± 3152.90	3428.70 ± 2780.22(n = 23)	−0.934(.356)
Depression	8.00 ± 2.77 (0–21)	7.16 ± 3.27 (0–21)	0.954 (.345)
Normal (≤7)	10(43.5)	14(56.0)	
Borderline abnormal (8–10)	9(39.1)	7(28.0)	
Abnormal (11–21)	4(17.4)	4(16.0)	
Anxiety	7.50 ± 3.57 (0–21)	7.88 ± 3.19 (0–21)	−0.380(.705)
Normal (≤7)	10(43.5)	14(56.0)	
Borderline abnormal (8–10)	9(39.1)	7(28.0)	
Abnormal (11–21)	4(17.4)	4(16.0)	
Sleep quality	8.64 ± 3.26 (0–21)	9.70 ± 3.55 (0–21)	−1.043(.302)
(Int: n = 21, Cont: n = 25)			
Normal (<8)	7(30.4)	6(24.0)	
Sleep disorder (≥8)	14(60.9)	19(76.0)	
Missing data	2(8.7)	0(0.0)	
Social capital	2.90 ± 0.63 (0–5)	2.80 ± 0.65 (0–5)	0.522(.604)
Bonding capital	3.40 ± 0.68	3.19 ± 0.76	1.011(.317)
Bridging capital	2.40 ± 0.72	2.42 ± 0.85	−0.077(.939)

Table 3 Comparison of dependent variables between groups (N = 48)

Variables	Group	Pretest Mean ± SD	Posttest Mean ± SD	Difference Mean ± SD	t/z	p
Fatigue	Int	5.24 ± 1.53	3.89 ± 1.19	1.34 ± 1.50	1.992	.052
	Con	5.37 ± 1.80	4.88 ± 1.52	0.48 ± 1.50		
Behavioral/Severity	Int	5.55 ± 1.42	3.87 ± 1.43	1.68 ± 1.57	2642	.011
	Con	5.46 ± 2.26	5.02 ± 2.11	0.43 ± 1.67		
Affective	Int	5.84 ± 2.11	4.86 ± 1.94	0.97 ± 1.89	1.674	.101
	Con	5.89 ± 2.55	5.86 ± 2.14	0.03 ± 2.02		
Sensory	Int	4.70 ± 2.31	3.57 ± 1.84	4.52 ± 9.52	0.571	.571
	Con	5.53 ± 1.87	4.79 ± 2.06	2.96 ± 9.41		
Cognitive/mood	Int	4.80 ± 2.00	3.39 ± 1.50	7.04 ± 9.35	1.331	.190
	Con	4.72 ± 2.05	4.03 ± 1.66	3.44 ± 9.38		
Quality of life	Int	85.58 ± 11.86	118.43 ± 17.63	-32.85 ± 15.10	-0.986	.329
	Con	86.80 ± 20.31	115.20 ± 24.95	-28.40 ± 16.10		
Physical wellbeing	Int	17.73 ± 4.70	20.69 ± 4.20	-2.95 ± 4.82	-0.584	.562
	Con	17.12 ± 6.48	19.32 ± 6.13	-2.20 ± 4.14		
Social/family wellbeing	Int	15.88 ± 5.46	16.81 ± 4.27	-0.92 ± 3.74	-1.358	.181
	Con	17.76 ± 5.20	17.24 ± 5.39	0.51 ± 3.64		
Emotional wellbeing	Int	15.26 ± 3.55	41.70 ± 8.33	-26.44 ± 7.94	-1.254	.216
	Con	15.36 ± 4.59	38.32 ± 11.89	-22.96 ± 10.91		
Functional wellbeing	Int	14.86 ± 4.54	16.56 ± 3.70	-1.69 ± 4.66	-0.395	.695
	Con	16.28 ± 4.79	17.48 ± 4.28	-1.20 ± 4.03		
Breast cancer subscale	Int	21.82 ± 4.31	22.65 ± 4.40	-0.83 ± 4.35	1.501	.140
	Con	20.28 ± 5.38	22.84 ± 4.91	-2.56 ± 3.61		
Physical activity MET, (n = 23/23)	Int	2610.43 ± 3152.90	4849.57 ± 3139.27	-2239.13 ± 4000.97	-2.049	.046
	Con	3428.70 ± 2780.22	3332.56 ± 2323.68	-130.17 ± 2889.58		
Depression	Int	8.00 ± 2.77	6.95 ± 2.91	1.04 ± 2.36	0.699	.488
	Con	7.16 ± 3.27	6.64 ± 2.41	0.52 ± 2.78		
Anxiety	Int	7.50 ± 3.57	7.04 ± 3.44	0.46 ± 3.04	-0.634	.529
	Con	7.88 ± 3.19	6.93 ± 3.00	0.94 ± 2.18		
Sleep quality	Int	8.64 ± 3.26	9.17 ± 3.47	-0.59 ± 2.93	-.648	.521
	Con	9.70 ± 3.55	9.60 ± 4.07	-0.04 ± 2.72		
Social capital	Int	2.90 ± 0.63	2.92 ± 0.64	-0.02 ± 0.48	0.493	.625
	Con	2.80 ± 0.65	2.89 ± 0.47	-0.09 ± 0.43		
Bonding capital	Int	3.40 ± 0.68	3.33 ± 0.67	0.07 ± 0.48	1.469	.149
	Con	3.19 ± 0.76	3.32 ± 0.54	-0.13 ± 0.46		
Bridging capital	Int	2.40 ± 0.72	2.52 ± 0.83	-0.11 ± 0.66	-0.330	.743
	Con	2.42 ± 0.85	2.47 ± 0.60	-0.05 ± 0.78		