



Correction to: The Role of Aprepitant in Prevention of Postoperative Nausea and Vomiting After Bariatric Surgery

Isaac W. Therneau¹ · Erin E. Martin¹ · Juraj Sprung¹ · Todd A. Kellogg² · Darrell R. Schroeder³ · Toby N. Weingarten¹

Published online: 11 February 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to: OBES SURG (2018) 28:37–43
<https://doi.org/10.1007/s11695-017-2797-0>

In the section “Discussion” on page 41, the second sentence in the second paragraph in the right-hand column should read as follows:

A reduction in PONV may be expected with aggressive antiemetic prophylaxis, and indeed, the PONV rate in PACU was 11 and 17% in aprepitant versus no-aprepitant groups, which reflect the protective effect of our intense perioperative antiemetic prophylactic regimen.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11695-017-2797-0>

✉ Toby N. Weingarten
weingarten.toby@mayo.edu

¹ Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, 200 First St SW, Rochester, MN 55905, USA

² Division of Subspecialty General Surgery, Mayo Clinic, Rochester, MN, USA

³ Division of Biomedical Statistics and Informatics, Mayo Clinic, Rochester, MN, USA